



Diphtheria, Tetanus, Pertussis, Polio, *Haemophilus influenzae* Type b (DTaP-IPV-Hib) Vaccine

**Keep your child safe.
Get all shots on time.**

By getting all shots on time, your child can be protected from many diseases over a lifetime.

Immunization has saved more lives in Canada in the last 50 years than any other health measure.

What is the DTaP-IPV-Hib vaccine?

The DTaP-IPV-Hib vaccine protects against five diseases:

- Diphtheria
- Tetanus
- Pertussis (or whooping cough)
- Polio
- *Haemophilus influenzae* type b

The vaccine is approved by Health Canada and is provided free as part of your child's routine immunizations. Call your public health nurse or doctor to make an appointment.

Who should get the DTaP-IPV-Hib vaccine?

Infants in B.C. receive DTaP-IPV-Hib as a booster dose at 18 months of age after completing a 3 dose primary series of DTaP-HB-IPV-Hib, which also protects against hepatitis B infection.

If your baby has been started on a primary series of DTaP-IPV-Hib, they should complete their vaccinations with this vaccine. They should also receive the hepatitis B vaccine. These vaccines should be given together at 2, 4 and 6 months of age. Your baby should receive a booster dose of DTaP-IPV-Hib at 18 months of age.

For more information, see HealthLink BC Files [#105 Diphtheria, Tetanus, Pertussis, Hepatitis B, Polio, and Haemophilus Influenzae Type b \(DTaP-HB-IPV-Hib\) Vaccine](#) and [#25C Hepatitis B Infant Vaccine](#).

It is important to keep a record of all immunizations received.

Benefits of DTaP-IPV-Hib Vaccine

The DTaP-IPV-Hib vaccine protects your child against diphtheria, tetanus, pertussis, polio, and *Haemophilus influenzae* type b, which are serious and sometimes fatal diseases.

When you get your child vaccinated, you help protect others as well.

Possible Reactions after the Vaccine

Vaccines are very safe. It is much safer to get the vaccine than to get the disease.

Common reactions to the vaccine may include soreness, redness and swelling where the vaccine was given. Some children may have a fever, or experience crankiness, drowsiness, or a loss of appetite. These reactions are mild and generally last 1 to 2 days. Larger areas of redness and swelling are more common after the booster dose at 18 months of age. These generally do not interfere with normal activity.

Acetaminophen or Tylenol® can be given for fever or soreness. ASA or Aspirin® should NOT be given to anyone under 20 years of age due to the risk of Reye Syndrome.

It is important to stay in the clinic for 15 minutes after getting any vaccine because there is an extremely rare possibility of a life-threatening allergic reaction called anaphylaxis. This may include hives, difficulty breathing, or swelling of the throat, tongue or lips. If this happens after you leave the clinic, call 911 or the local emergency number. This reaction can be treated and occurs in less than one in a million people who get the vaccine.

Report serious or unexpected reactions to your public health nurse or doctor.

Who should not get the DTaP-IPV-Hib vaccine?

Speak with a public health nurse or doctor if your child has had a life-threatening reaction to a previous dose of diphtheria, pertussis, tetanus, polio, or Hib vaccine, or any component of the vaccine, including neomycin or polymyxin B. The vaccine is not given after 5 years of age.

People who developed Guillain-Barré Syndrome (GBS) within 8 weeks of getting tetanus vaccine should not get the DTaP-IPV-Hib vaccine. GBS is a rare condition that can result in weakness and paralysis of the body's muscles.

What are Diphtheria, Pertussis, Tetanus, Polio, and *Haemophilus influenzae* type b?

Diphtheria is a serious infection of the nose and throat. About 1 in 10 people who get diphtheria may die. The germ (bacteria) is spread through the air from person to person. The disease can result in very severe breathing problems. It can also cause heart failure and paralysis.

Pertussis, or “whooping cough”, is a serious infection of the lungs and throat. About 1 in 2000 people who get pertussis may die. Pertussis can cause pneumonia, convulsions, brain damage or death. These complications are seen most often in infants. The germ (bacteria) is easily spread by coughing, sneezing or close face-to-face contact. Pertussis can cause severe coughing that often ends with a whooping sound before the next breath. This cough can last several months and occurs more often at night.

Tetanus, also called “lockjaw”, is caused by a germ (bacteria) mostly found in the soil. When the bacteria enter the skin through a cut or scrape, they produce a poison that can cause painful tightening of muscles all over the body. It is very serious if the breathing muscles are affected. Up to 2 in 10 people who get tetanus may die.

Polio is a disease caused by infection with a virus. While some polio infections show no symptoms, others can result in paralysis of arms or legs and even death. Polio can be spread by contact with the bowel movements (stool) of an infected person. This can happen from eating food or drinking water contaminated with stool.

***Haemophilus influenzae* type b** (Hib) is a germ (bacteria) that most commonly infects children under 5 years old. It can cause serious and life-threatening infections including meningitis, an infection of the lining that covers the brain, and septicemia, an

infection of the blood. For every 20 children who get sick, one may die. Hib infection is spread by coughing, sneezing or having close face-to-face contact.

Four of these diseases are now rare in B.C. because of routine childhood vaccination programs. Whooping cough still occurs, but it is far less common than it used to be and it is much milder in immunized people.

For more HealthLink BC File topics, visit www.HealthLinkBC.ca/healthfiles/index.stm or your local public health unit.

Click on www.HealthLinkBC.ca or call 8-1-1 for non-emergency health information and services in B.C.

For deaf and hearing-impaired assistance, call 7-1-1 in B.C.

Translation services are available in more than 130 languages on request.



ImmunizeBC



BC Centre for Disease Control
AN AGENCY OF THE PROVINCIAL HEALTH SERVICES AUTHORITY