

E. coli infection

What is *E. coli* infection?

E. coli infection is a type of diarrheal illness caused by bacteria called *Escherichia coli*. It affects the digestive tract and in serious cases, the kidneys. There are many strains of *E. coli* and most are harmless.

When food or water tests positive for *E. coli* bacteria, it may be an indication of fecal contamination.

The strain of *E. coli* that most often makes people sick is Shiga-toxin producing *E. coli*. For ease, we will refer to Shiga-toxin producing *E. coli* as *E. coli*.

What are the symptoms?

Symptoms of *E. coli* infection include:

- Watery diarrhea, which in serious cases may become bloody
- Nausea
- Vomiting
- Stomach cramps
- Mild fever

The symptoms can start 2 to 10 days after contact with or consuming something that contains *E. coli*, but usually starts within 3 to 4 days. Symptoms can last for 5 to 10 days.

In some cases, *E. coli* can cause a serious and sometimes fatal illness called hemolytic uremic syndrome (HUS) which can result in kidney failure, anemia and internal bleeding. HUS can be especially harmful for young children, the elderly, people with weakened immune systems and pregnant women.

Where does the disease come from?

E. coli is present in the intestines and feces of many healthy animals including cattle, goats, sheep, deer and elk.

During the butchering process, *E. coli* sometimes gets onto the surface of the meat. Whole cuts of

meat such as steaks or roasts usually only have *E. coli* on the surface, which makes the *E. coli* easier to kill by cooking.

When the meat is ground or mechanically tenderized, *E. coli* on the surface can be transferred to the inside of the meat. This is why ground meat and mechanically tenderized meat are more likely to cause illness than whole cuts of meat. *E. coli* can be killed if the meat is cooked thoroughly.

E. coli are also sometimes found in other foods including fruits and vegetables, as well as in unpasteurized milk products, juice, cider and untreated or contaminated drinking water.

Swimming in recreational water that is contaminated with human or animal sewage can also cause infections.

E. coli can also be transmitted from a sick person to another person.

How do I avoid getting sick?

To reduce the risk of *E. coli* infection:

- Refrigerate or freeze meat as soon as possible after buying it
- Always thaw meat thoroughly in the refrigerator before you start to cook it. Ensure it is wrapped well and keep away from other foods. Do not try cooking frozen or partially frozen meat
- Wash your hands thoroughly before preparing food, after handling raw meat and after using the toilet or changing diapers as *E. coli* can be spread to others through infected feces
- Always wash and then sanitize all utensils, cutting boards and counters that have touched raw meat to prevent *E. coli* from contaminating other foods. You can use a 200 ppm (parts per million) no rinse sanitizing solution

How to make a 200 ppm no rinse sanitizing solution:

1. Mix 15 mL (1 tablespoon) of household bleach into 4 litres (1 gallon) of water; or mix 5 mL (1 teaspoon) of household bleach into 1 litre (4 cups) of water
2. Allow the sanitizer to contact the surface or utensil for at least 1 minute before wiping off with a clean paper towel or allowing to air dry

Use B.C.'s FOODSAFE Chlorine Dilution Calculator tool to make up the proper sanitizer strength based on the concentration of your bleach product
www.foodsafe.ca/dilution-calculator.html

Caution: Mixing bleach with products that contain acid or ammonia makes a toxic gas that causes serious breathing problems, choking and potentially death.

- Cook whole cuts of beef to a minimum internal temperature of 63°C (145°F). If cooking steaks, turn at least twice during cooking
- Cook ground meat, such as hamburger patties, meat loaf and rolled roasts (beef roulade) and mechanically tenderized cuts of beef until the middle reaches a temperature of 74°C (164°F) or hotter and remains there for at least 15 seconds. Measure the internal temperature of the meat with a digital probe tip thermometer
- Never put cooked meat back on the same unwashed surface that held the raw meat. The juices from the raw meat can be contaminated with *E. coli*, which can then contaminate the cooked meat
- Do not re-use marinade as a sauce on cooked meat unless the marinade is boiled first
- Serve cooked meats right away or keep them hot. They should be at least 60°C (140°F) or hotter, until you are ready to serve them
- Wash any fruits and vegetables with cold running water before eating them raw, even if they will be peeled or cut. Use a scrub brush on

fruits and vegetables with rough skin, such as a cantaloupe

- Avoid drinking unpasteurized milk, juice and cider
- Do not drink or use untreated surface water from a spring, stream, river, lake, pond or shallow well. Assume it is contaminated with animal feces
- Avoid swimming in closed beaches or water that may drain pastureland

In B.C., drinking water from the tap is routinely tested for generic *E. coli* contamination.

Is there a treatment for the disease?

If you have diarrhea and/or bloody diarrhea for more than a few days, you should see a health care provider.

Drinking lots of fluids is important to replace lost fluids and prevent dehydration. Do not take anti-diarrhea medicine or antibiotics unless recommended by your health care provider.

If you have a serious *E. coli* infection, you may need hospitalization, blood transfusion and dialysis.

For more information

For more information about food safety, see the following HealthLinkBC Files:

- [HealthLinkBC File #03 Pasteurized and raw milk](#)
- [HealthLinkBC File #22 Home canning - How to avoid botulism](#)
- [HealthLinkBC File #59a Food safety: Easy ways to make food safe](#)
- [HealthLinkBC File #59d Food safety in child care facilities](#)
- [HealthLinkBC File #72 Unpasteurized fruit juices and ciders: A potential health risk](#)



BC Centre for Disease Control
Provincial Health Services Authority