



Grade 9 immunizations in B.C.

Immunization has saved more lives in Canada in the last 50 years than any other health measure.

Vaccines are the best way to protect your child against many diseases and their complications. When you get your child immunized, you help protect others as well.

The meningococcal quadrivalent conjugate vaccine and Tdap vaccine are offered to all students in grade 9. Your child may be offered other vaccines if they have not had all doses of the recommended childhood vaccines that they are eligible for. For more information on the recommended childhood vaccines in B.C., see the B.C. Immunization Schedules at <u>www.healthlinkbc.ca/health-</u> <u>library/immunizations/schedules</u>.

It is important to keep a record of all immunizations that your child has received. If your child received any vaccines outside of B.C., you can update their immunization record at

www.immunizationrecord.gov.bc.ca.

What is the meningococcal quadrivalent conjugate vaccine?

The meningococcal quadrivalent conjugate vaccine protects against infection from 4 types of meningococcal bacteria: types A, C, Y and W. It replaced the meningococcal C (Men-C) vaccine that was previously offered to students in grade 6. Children should get the meningococcal quadrivalent conjugate vaccine in grade 9 even if they received the Men-C vaccine when they were younger.

For more information, see <u>HealthLinkBC File #23b</u> <u>Meningococcal Quadrivalent Vaccines</u>.

What is meningococcal infection?

Meningococcal infection is caused by bacteria. Meningococcal infection due to types A, C, Y and W is very rare in B.C. Although rare, it can cause serious and life-threatening infections. These include meningitis, an infection of the lining that covers the brain, and septicemia, an infection of the blood. For every 100 people who get sick, up to 15 will die, even if treated. Permanent complications include brain damage, deafness and loss of limbs.

Meningococcal infection spreads by coughing, sneezing or close face-to-face contact. It may spread through saliva by kissing, or sharing of food, drinks, cigarettes, lipstick, water bottles, mouth guards used for sports and mouthpieces of musical instruments.

What is the Tdap vaccine?

The Tdap vaccine protects against 3 diseases:

- Tetanus
- Diphtheria
- Pertussis (or whooping cough)

The Tdap vaccine offered in grade 9 is a booster dose for children who were immunized against these diseases at a younger age. The booster dose strengthens or boosts the immune system to give better protection against these diseases. If your child had a dose of Tdap vaccine on or after their 10th birthday, they do not need a dose in grade 9.

Also see <u>HealthLinkBC File #18c Tetanus, Diphtheria,</u> <u>Pertussis (Tdap) Vaccine</u>.

What are tetanus, diphtheria & pertussis?

Tetanus, also known as lockjaw, is caused by bacteria mostly found in soil. When the bacteria enter the skin through a cut or scrape, they produce a poison that can cause painful tightening of muscles all over the body. It is very serious if the breathing muscles are affected. Up to 1 in 5 people who get tetanus may die.

Diphtheria is a serious infection of the nose and throat caused by diphtheria bacteria. The bacteria spread through the air when people sneeze or cough, or by direct skin-to-skin contact. The disease can result in severe breathing problems. It can also cause heart failure and paralysis. About 1 in 10 people who get diphtheria may die. **Pertussis**, also known as whooping cough, is a serious infection of the airways caused by pertussis bacteria. Pertussis can cause pneumonia, seizures, brain damage or death. These complications are seen most often in infants. The bacteria spreads easily when someone coughs, sneezes, or has close face-to-face contact with another person. Pertussis can cause severe coughing that often ends with a whooping sound before the next breath. This cough can last several months and occurs more often at night. About 1 in 170 infants who get pertussis may die.

What are the possible reactions after these vaccines?

Common reactions may include soreness, redness and swelling in the arm where the vaccine was given. You may also have headache, fatigue, muscle or joint soreness, nausea, irritability, loss of appetite, chills, and fever.

Acetaminophen (e.g. Tylenol[®]) or ibuprofen (e.g. Advil[®]) can be given for fever or soreness. ASA (e.g. Aspirin[®]) should not be given to anyone under 18 years of age due to the risk of Reye Syndrome.

For more information on Reye Syndrome, see <u>HealthLinkBC File #84 Reye syndrome</u>.

It is important to stay in the clinic for 15 minutes after getting any immunization because there is an extremely rare possibility, less than 1 in a million, of a life-threatening allergic reaction called anaphylaxis. This may include hives, difficulty breathing, or swelling of the throat, tongue or lips. If this reaction occurs, your health care provider is prepared to treat it. Emergency treatment includes administration of epinephrine (adrenaline) and transfer by ambulance to the nearest emergency department. If symptoms develop after you leave the clinic, call **9-1-1** or the local emergency number.

It is important to always report all serious or unexpected reactions to your health care provider.

Who should not get a vaccine?

A vaccine is not recommended for:

- People who have had a life-threatening allergic reaction to a previous dose of vaccine, or to any part of the vaccine
- People who developed Guillain-Barré Syndrome (GBS) within 8 weeks of getting a tetanus vaccine, without another cause being identified, should not get the Tdap vaccine. GBS is rare and can result in weakness and paralysis of the body's muscles. It most commonly occurs after infections. In rare cases, GBS can also occur after some vaccines

There is no need to delay getting immunized because of a cold or other mild illness. However, if you have concerns, speak with your health care provider.

Mature minor consent

It is recommended that parents or guardians and their children discuss consent for immunization. Children under the age of 19, who are able to understand the benefits and possible reactions for each vaccine and the risk of not getting immunized, can legally consent to or refuse immunizations. For more information on mature minor consent see <u>HealthLinkBC File #119 The Infants Act, Mature</u> <u>Minor Consent and Immunization</u>.

BC Centre for Disease Control

For more HealthLinkBC File topics, visit <u>www.HealthLinkBC.ca/more/resources/healthlink-bc-files</u> or your local public health unit. For non-emergency health information and advice in B.C. visit <u>www.HealthLinkBC.ca</u> or call **8-1-1** (toll-free). For the deaf and hard of hearing, call **7-1-1**. Translation services are available in more than 130 languages on request.