

## Immunization Schedule for B.C. Infants and Children

Vaccine	2 Months	4 Months	6 Months	12 Months	18 Months	Starting at 4 Years of Age (Kindergarten Entry)
<b>Chickenpox (Varicella) Vaccine</b> (#44b) <sup>1</sup>				<input checked="" type="checkbox"/>		
<b>Diphtheria, Tetanus, Pertussis, Hepatitis B, Polio, and Haemophilus influenzae type b (DTaP-HB-IPV-Hib) Vaccine</b> (#105)	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>			
<b>Diphtheria, Tetanus, Pertussis, Polio, Haemophilus influenzae Type b (DTaP-IPV-Hib) Vaccine</b> (#15b)					<input checked="" type="checkbox"/>	
<b>Hepatitis A Vaccine</b> (#33) Indigenous children only			<input checked="" type="checkbox"/>		<input checked="" type="checkbox"/>	
<b>Inactivated Influenza (Flu) Vaccine</b> (#12d) <sup>2</sup>			<input checked="" type="checkbox"/> Annually for children 6 months to 4 years of age			
<b>Measles, Mumps, Rubella (MMR) Vaccine</b> (#14a)				<input checked="" type="checkbox"/>		
<b>Measles, Mumps, Rubella and Varicella (MMRV) Vaccine</b> (#14e) <sup>1</sup>						<input checked="" type="checkbox"/>
<b>Meningococcal C Conjugate (Men-C) Vaccine</b> (#23a)	<input checked="" type="checkbox"/>			<input checked="" type="checkbox"/>		
<b>Pneumococcal Conjugate (PCV 13) Vaccine</b> (#62a)	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>		<input checked="" type="checkbox"/>		
<b>Rotavirus Vaccine (RotaTeq®)</b> (#104)	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>			
<b>Tetanus, Diphtheria, Pertussis, Polio (Tdap-IPV) Vaccine</b> (#15a)						<input checked="" type="checkbox"/>

**Note:** The vaccine schedule can change. Speak with your health care provider, or call **8-1-1** if you have questions. This immunization table was developed and reviewed by HealthLinkBC, BC Ministry of Health and BC Centre for Disease Control.

<sup>2</sup> Annual influenza immunization is recommended and provided free to children 6 months to 4 years of age. For children receiving an influenza vaccine for the first time, a second dose is recommended 4 weeks after the first dose.

<sup>1</sup> Children who have had chickenpox or shingles disease, confirmed by a lab test, at 1 year of age or older do not need the chickenpox vaccine, and should get the MMR vaccine at 4 years of age.