

Immunization Schedule for B.C. Infants and Children

Vaccine	2 Months	4 Months	6 Months	12 Months	18 Months	Starting at 4 Years of Age (Kindergarten Entry)
Chickenpox (Varicella) Vaccine (#44b) ¹				<input checked="" type="checkbox"/>		
Diphtheria, Tetanus, Pertussis, Hepatitis B, Polio, and Haemophilus influenzae type b (DTaP-HB-IPV-Hib) Vaccine (#105)	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>			
Diphtheria, Tetanus, Pertussis, Polio, Haemophilus influenzae Type b (DTaP-IPV-Hib) Vaccine (#15b)					<input checked="" type="checkbox"/>	
Hepatitis A Vaccine (#33) Aboriginal children only			<input checked="" type="checkbox"/>		<input checked="" type="checkbox"/>	
Influenza Vaccine Inactivated Influenza (Flu) Vaccine (#12d) ² Live Attenuated Influenza (Flu) Vaccine (#12e) ²			<input checked="" type="checkbox"/> Children 6 months to 4 years of age (The live attenuated influenza vaccine is not approved for use in children under 2 years of age)			
Measles, Mumps, Rubella (MMR) Vaccine (#14a)				<input checked="" type="checkbox"/>		
Measles, Mumps, Rubella and Varicella (MMRV) Vaccine (#14e) ¹						<input checked="" type="checkbox"/>
Meningococcal C Conjugate (Men-C) Vaccine (#23a)	<input checked="" type="checkbox"/>			<input checked="" type="checkbox"/>		
Pneumococcal Conjugate (PCV 13) Vaccine (#62a)	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>		<input checked="" type="checkbox"/>		
Rotavirus Vaccine (RotaTeq®) (#104)	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>	<input checked="" type="checkbox"/>			
Tetanus, Diphtheria, Pertussis, Polio (Tdap-IPV) Vaccine (#15a)						<input checked="" type="checkbox"/>

Note: The vaccine schedule can change. Speak with your health care provider, or call **8-1-1** if you have questions. This immunization table was developed and reviewed by HealthLinkBC, BC Ministry of Health and BC Centre for Disease Control.

² Annual influenza immunization is recommended and provided free to children 6 months to 4 years of age. For children receiving an influenza vaccine for the first time, a second dose is recommended 4 weeks after the first dose.

¹ Children who have had chickenpox or shingles disease, confirmed by a lab test, at 1 year of age or older do not need the chickenpox vaccine, and should get the MMR vaccine at 4 years of age.