



Clostridium Difficile (*C.difficile*)

What is *C.difficile*?

C. difficile, also known as *C.diff*, is a bacteria that lives in the intestines of 1 to 3% of people. Your intestines also normally contain many good bacteria that help you digest food and stay healthy. When antibiotics are taken to treat an illness, these good bacteria can be killed. *C.diff* bacteria can survive and cause you to become sick.

C. diff produces toxins that can cause damage to the cells in the intestines. The most common symptom of *C.diff* is diarrhea. In fact, it is the most frequent cause of infectious diarrhea in hospitals and health care facilities. The reported number of healthcare associated *C. diff* infections in Canada has risen over the last decade in hospitals, and is linked with an increase in illnesses and sometimes death.

Who is most at risk?

For most healthy people, *C. diff* is not a health risk. Those at greater risk of infection include:

- people who are taking antibiotics;
- people who have other illnesses; and
- the elderly.

What are the symptoms?

In some cases *C. diff* may cause no symptoms. However, it may also cause symptoms such as:

- watery diarrhea;
- fever;
- loss of appetite;
- nausea; and
- abdominal pain and tenderness.

In cases where a patient has severe diarrhea, it can lead to complications including dehydration and colitis (inflammation of the lower intestine or colon). In rare cases, the infection can cause death.

How is *C. diff* spread?

C.diff bacteria and their spores are found in feces. The spores can live outside your body for weeks or months. They may be found on items you touch such as bed linens, bed rails, bathroom fixtures, and medical equipment.

You can become infected if you touch a surface contaminated with feces and then touch your mouth, or if you eat or drink something that is contaminated.

Healthcare workers, staff and visitors can spread the infection to you if their hands have come in contact with the *C.diff* bacteria, and they have not properly cleaned their hands before touching you, or touching items that you might eat or drink.

C. diff bacteria are not spread through the air. You cannot get *C. diff* by someone sneezing or coughing.

How can *C. diff* be prevented?

The best way to stop the spread of infection is to regularly wash your hands with soap and water, especially after using the toilet and before you eat.

If you develop symptoms, such as diarrhea, that are either confirmed or suspected to be from *C. diff*, you will be moved to a room by yourself. You will also be put on contact precautions to try to prevent spreading the infection to others.

How can you wash your hands?

To wash your hands properly, follow the steps below:

1. Remove rings or other jewelry on your hands and wrists.
2. Wet your hands with warm water.
3. Wash all parts of your hands with soap and water for at least 20 seconds, and rub your hands together to create lather. To help children wash their hands, sing the alphabet song.
4. Rinse hands well under warm running water.
5. Dry your hands with a clean cloth or paper towel.

6. Use the towel to turn off the tap and open the door when you leave if you are in a public restroom.

If washing hands frequently apply hand cream often to prevent skin breakdown.

For more information on hand washing, see [HealthLinkBC File #85 Hand Washing for Parents and Children](#).

What are contact precautions?

Contact precautions are actions taken by health care facilities to try to prevent the spread of *C. diff* bacteria to other patients.

Contact precautions include placing all patients with suspected and confirmed cases of *C. diff* into single isolation rooms. If single rooms are not available, infected patients can share a room if they have the same type of *C. diff* bacteria. Doors must be kept closed and a “Contact Precautions” sign placed on the door.

Before entering the room of a patient who has *C. diff* the following actions must be taken by health care providers, staff and visitors:

1. Wash your hands.
2. Put an isolation gown on over your clothing. This will help to avoid contact with any contaminated surfaces or with the patient.
3. Put on non-sterile gloves.

When leaving a patient’s room, the following actions must be taken:

1. Remove gloves immediately and throw them in the garbage.
2. Wash your hands.
3. Remove isolation gown without touching the outside of the gown and place it in the linen hamper provided.
4. Wash your hands for a second time.
5. Exit the room by opening the door with a paper towel.

6. Wash your hands for a third time. The inside of the door handle may have been contaminated. Health care facilities will also frequently clean surfaces with sporacidal bleach to reduce the number of bacteria.

What are the treatments?

The first step in treating a *C. diff* infection is to stop taking the antibiotic that caused the infection. However, you should not stop taking your antibiotics until you have been told to do so by your health care provider. Although antibiotics can cause *C. diff*, there are also some antibiotics that can be used to treat *C. diff*. Your health care provider will prescribe an antibiotic to treat your *C. diff* infection.

In rare cases, surgery may be considered if all other treatments have not worked, or the symptoms are very severe.



BC Centre for Disease Control
An agency of the Provincial Health Services Authority

For more HealthLinkBC File topics, visit www.HealthLinkBC.ca/healthlinkbcfiles or your local public health unit. For non-emergency health information and advice in B.C. visit www.HealthLinkBC.ca or call **8-1-1** (toll-free). For deaf and hearing-impaired assistance, call **7-1-1**. Translation services are available in more than 130 languages on request.