



HTLV – 1 Virus

What is HTLV-1?

Human T-Lymphotropic Virus Type 1 (HTLV -1) is a virus that infects human white blood cells. It can cause disease of the nervous system or leukemia.

Is this a new disease?

No. This disease was first described in 1980 and has been identified throughout the world.

HTLV-1 is more common in Japan and other countries in the Western Pacific region, the Caribbean, West Africa and South America. About 10 to 20 million people are infected worldwide. It is rare in Canada, but cases have been found in British Columbia and Nunavut.

How can I tell if I have HTLV-1?

The majority of people who have the virus will not get symptoms or develop any health problems from this virus.

About 1 in 20 people who get infected by HTLV-1 will eventually get sick with an HTLV-1-associated illness sometime in their lifetime. Illness does not usually occur until several decades after being infected.

People who get sick may develop a rare blood cancer called Adult T-cell leukemia/lymphoma (ATLL), or have inflammation of the spinal cord that causes weakness of the legs, backache, loss of bladder control or constipation.

To confirm that you have HTLV-1, your health care provider may send you for a blood test.

How is the virus spread?

HTLV-1 is spread from an infected person to other people by:

- Sharing needles, syringes or other equipment used for injecting drugs.
- Sexual contact. Evidence shows that the virus is spread more easily from a man to a woman, than from a woman to a man. The virus is also more common in people who are 40 years or older.
- Mother to child. About one quarter of mothers who are infected with HTLV-1 may transmit the virus to their babies at birth or through breastfeeding, especially if they breastfeed for 6 months or more.

Should I breastfeed if I have HTLV-1?

If you are infected with HTLV-1 you should not breastfeed. Otherwise, breastfeeding is highly recommended. For more information on breastfeeding, see [HealthLinkBC File #70 Breastfeeding](#).

How can I protect myself against infection from HTLV-1?

You can protect yourself from being infected with HTLV-1 by:

- Never sharing needles, syringes or other drug injection equipment.
- Using a condom every time you have vaginal, anal or oral sex.

HTLV-1 is not spread by ordinary, everyday household contacts such as kissing, using the toilet or preparing food. In order for the disease to spread, there has to be blood-to-blood contact or unprotected sexual intercourse.

Is there any treatment for HTLV-1 infection?

No. There is currently no treatment that will get rid of the virus once you are infected, and 95 percent of those who are infected do not develop any associated illness. However, some treatments exist for illnesses, such as ATLL, that may develop as a result of HTLV-1 infection.

Can I get HTLV-1 from blood transfusions?

Canadian Blood Services (formerly Canadian Red Cross) has been screening all blood donations for HTLV-1 since April 1990 and transmission of HTLV-1 via transfusion has been virtually eliminated in Canada.

Should I get tested?

No, unless you have a HTLV-1-associated illness for which your health care provider thinks testing is needed, or you have been in contact with a known case.

For more HealthLinkBC File topics, visit www.HealthLinkBC.ca/healthfiles or your local public health unit.

Click on www.HealthLinkBC.ca or call **8-1-1** for non-emergency health information and services in B.C.

For deaf and hearing-impaired assistance, call 7-1-1 in B.C.

Translation services are available in more than 130 languages on request.



BC Centre for Disease Control
An agency of the Provincial Health Services Authority